INTERNATIONAL JOURNAL OF MACHINE LEARNING AND ENGINEERING

Vol 2 Issue 1 (2026)

https://ijmle.org/

Calibrating Fatigue Scales in Autoimmune Trials Across Multicenter Autoimmune Patient Cohorts

Joao Campos, Akane Miyazaki, Gavin Parker

Faculty of Science, University of Helsinki, Helsinki, Finland

Abstract:

The accurate measurement of fatigue in multicenter autoimmune disease trials is a critical challenge due to the subjective nature of the symptom and the diverse cultural backgrounds of trial participants. Patient-reported outcomes have become central to evaluating therapeutic efficacy in conditions such as rheumatoid arthritis, systemic lupus erythematosus, and multiple sclerosis. However, the utilization of standard fatigue scales across multinational cohorts frequently introduces measurement variance, particularly differential item functioning, where demographic and linguistic differences distort symptom reporting. This paper presents a comprehensive analysis of the methodological frameworks required for the psychometric calibration of patient-reported fatigue scales. Transitioning from classical test theory to modern approaches based on Item Response Theory and Rasch measurement, we detail the procedures necessary to establish measurement equivalence across diverse geographic centers. Through simulated trial data, we demonstrate how uncalibrated ordinal scores obscure true treatment effects and how advanced calibration techniques, including item splitting and concurrent anchoring, neutralize cultural bias. The results highlight that scale calibration significantly improves statistical power and ensures that the clinical trial data are fit for regulatory evaluation. Ultimately, robust calibration methodologies are indispensable for maximizing the validity of subjective endpoints in global clinical research.

Keywords: *Patient-reported outcomes, Fatigue scales, Autoimmune diseases, Calibration.*

INTRODUCTION

1.1 The Role of Patient-Reported Outcomes

Autoimmune diseases present a vast array of systemic symptoms, among which fatigue is consistently reported by patients as one of the most debilitating and pervasive issues. Fatigue in autoimmune conditions, particularly in rheumatoid arthritis, systemic lupus erythematosus, and multiple sclerosis, transcends ordinary tiredness, manifesting as a profound, unpredictable exhaustion that significantly impairs the overall quality of life [1]. Despite its high prevalence and severe impact on daily functioning, fatigue remains an inherently subjective experience, making it exceptionally challenging to quantify accurately in rigorous clinical research settings [2].

Historically, clinical trials heavily favored objective biological markers, but as pharmaceutical interventions increasingly target holistic symptom relief alongside fundamental disease modification, patient-reported outcomes have become critical primary and secondary endpoints [3]. Regulatory agencies globally, including those in North America and Europe, explicitly recognize the overarching importance of these subjective measures in determining the true clinical benefit of a novel therapeutic agent [4]. However, these regulatory bodies stipulate that any instrument used to capture patient-reported outcomes requires robust psychometric validation to ensure that the tools accurately capture the intended construct without being confounded by external or irrelevant variables [5]. The subjective nature of fatigue means that its measurement is highly susceptible to contextual influences, necessitating sophisticated analytical strategies to isolate the physiological burden of the disease from psychosocial noise.

1.2 Challenges in Multicenter Environments

In the specific context of multicenter clinical trials, the measurement of fatigue introduces substantial methodological complexities that can threaten the integrity of the entire study. Such large-scale trials often span multiple continents, encompassing highly diverse cultural, linguistic, and socio-economic patient populations [6]. When standard fatigue scales are translated and applied across these varying demographic groups, researchers frequently encounter differential item functioning, a psychometric phenomenon where individuals from different groups with the exact same underlying level of fatigue respond differently to specific scale items [7]. This variance can stem from subtle linguistic nuances in translation, profound cultural differences in the willingness to report psychological distress, or varied societal interpretations of somatic symptoms [8]. If left completely unaddressed by trial statisticians, this measurement variance can obscure true treatment effects, leading to either false positive or false negative efficacy results, thereby jeopardizing the massive financial and temporal investments required for a clinical development program [9]. Calibration of patient-reported fatigue scales is therefore an indispensable preliminary step in the design and subsequent analysis of multicenter autoimmune trials [10]. Calibration involves using advanced psychometric techniques to ensure that a survey scale measures the same latent construct with mathematically equivalent precision across all participating cohorts. By establishing strict measurement equivalence, clinical researchers can confidently pool data from different global centers, knowing unequivocally that a specific score on the fatigue scale signifies the identical level of symptom severity regardless of the geographic location or cultural background of the respondent.

2. Literature Review and Background

2.1 Evolution of Fatigue Measurement

The inclusion of fatigue as a primary or secondary endpoint in autoimmune disease trials has evolved significantly over the past three decades of clinical research. Early clinical evaluations primarily focused on objective, quantifiable biological markers, such as specific inflammatory cytokines, swollen joint counts, or radiological evidence of disease progression [11]. However, persistent discrepancies between objective disease activity metrics and patient-reported quality of life prompted a fundamental paradigm shift toward highly patient-centric outcome measures. The clinical realization that fatigue often operates entirely independently of traditional inflammatory markers in many chronic autoimmune conditions necessitated the immediate development of

specific survey instruments designed explicitly to capture this otherwise invisible symptom [12]. The Multidimensional Assessment of Fatigue, the Functional Assessment of Chronic Illness Therapy Fatigue scale, and the Fatigue Severity Scale eventually emerged as prominent and widely deployed tools in this endeavor, each positing a slightly different conceptual model of exhaustion. The Functional Assessment of Chronic Illness Therapy Fatigue scale has gained particular prominence in both rheumatology and oncology trials globally. Originally developed for cancer-related fatigue, its comprehensive thirteen-item structure evaluates both the physical limitations and social ramifications of extreme exhaustion. Countless validation studies have repeatedly confirmed its high internal consistency and test-retest reliability within specific, culturally homogenous populations [13].

2.2 Psychometric Limitations of Classical Theories

Despite the initial success of these instruments, their application expanded rapidly into global multicenter autoimmune trials, and researchers quickly began noting significant statistical inconsistencies in cross-cultural settings. Certain questionnaire items, particularly those closely related to social participation or the emotional burden of chronic fatigue, showed tremendous variance depending heavily on the cultural context of the respondents [14]. The limitations of classical test theory in evaluating and refining these scales have become increasingly apparent to the psychometric community. Classical test theory fundamentally assumes that a patient true score is simply the sum of observed scores and random measurement error, relying heavily on summary statistics that are entirely sample-dependent [15]. This mathematical dependency means that item difficulty and patient ability are not independently assessed or separated. Consequently, if a specific fatigue scale is validated exclusively in a predominantly Western European clinical cohort, its established psychometric properties may not hold true in an Asian, African, or Latin American patient population [16]. This sample dependency represents a critical methodological flaw when designing global phase three trials, where the most fundamental requirement is a highly stable metric that transcends all geographical and linguistic boundaries [17].

2.3 Modern Frameworks and Item Response Theory

To directly overcome these severe classical limitations, modern psychometrics has turned resolutely to Item Response Theory and Rasch measurement theory. These advanced mathematical frameworks offer a sophisticated paradigm where the statistical probability of a specific patient response is modeled as a mathematical function of both the respondent underlying trait level and the inherent properties of the survey item itself, such as its difficulty and discrimination capacity [18]. In Rasch analysis, the strict separation of item parameters and person parameters allows for the highly accurate construction of interval-level measurement from standard ordinal survey data, which is an absolutely crucial step for conducting valid parametric statistical analysis in clinical trials. The rigorous demands of the Rasch model, which absolutely requires unidimensionality and local independence of all included items, provide a robust, objective diagnostic tool for scale calibration [19]. If the trial data fit the Rasch model appropriately, researchers can be entirely confident that the simple sum score is a mathematically sufficient statistic for the latent trait of fatigue, completely irrespective of the sample distribution. The application of these modern psychometric techniques in multicenter trials has sharply highlighted the pervasive and troublesome issue of differential item functioning across global borders. Differential item

functioning occurs precisely when individuals with the exact same level of latent, underlying fatigue have differing mathematical probabilities of endorsing a specific item based merely on their group membership, such as their gender, age cohort, native language, or country of residence [20]. Identifying, quantifying, and mathematically adjusting for this differential functioning is the ultimate core objective of psychometric scale calibration.

3. Methodology and Analysis

3.1 Study Design and Qualitative Foundations

The comprehensive methodology for calibrating patient-reported fatigue scales in a multicenter clinical context requires a highly structured, multiphase approach that meticulously integrates both qualitative linguistic assessments and quantitative psychometric modeling. The initial phase of calibration involves establishing a highly robust data collection protocol that captures a sufficiently diverse patient sample to accurately represent the entire global trial population. In our analytical framework detailed herein, we rigorously evaluate simulated data derived directly from a hypothetical, large-scale multinational clinical trial encompassing patients formally diagnosed with systemic lupus erythematosus, rheumatoid arthritis, and multiple sclerosis. The sample must be carefully stratified to ensure adequate statistical representation across distinct geographic regions, primary language groups, and baseline disease severities. This stratification typically requires a minimum of two hundred respondents per specific demographic subgroup to achieve highly stable and reproducible parameter estimates in the subsequent Rasch analyses [21]. The qualitative foundation of calibration involves comprehensive linguistic validation and cognitive patient debriefing. Before any quantitative analysis can commence, clinical researchers must ensure beyond a doubt that questionnaire translations are not merely literal but conceptually equivalent across all languages. Forward and backward translation processes, immediately followed by rigorous expert panel reviews, help mitigate initial linguistic discrepancies. Cognitive debriefing interviews with a small, representative subset of patients in each target geographic region allow researchers to deeply probe how typical respondents interpret specific scale items and response categories [22].

Table 1: Baseline Characteristics of the Simulated Multicenter Autoimmune Cohort

Characteristic	European Cohort	Asian Cohort	North American Cohort
Total Participants	450	450	450
Mean Age in Years	47.5	45.2	48.1
Disease Duration in Years	8.3	7.9	9.1
Rheumatoid Arthritis Patients	150	150	150
Systemic Lupus Erythematosus Patients	150	150	150
Multiple Sclerosis Patients	150	150	150

Baseline Mean Fatigue Raw Score	38.4	41.2	39.7

3.2 Quantitative Calibration and Rasch Modeling

The quantitative calibration process employs the strict mathematical parameters of the Rasch measurement model to comprehensively evaluate scale performance across the pooled cohort. The first absolute requirement is assessing psychometric unidimensionality, which basically posits that all items in the chosen scale measure a single, isolated underlying construct, which in this specific case is disease-related fatigue. Principal component analysis of the model residuals is typically utilized to verify this fundamental assumption statistically. If the first residual contrast accounts for a variance equivalent to more than two individual items, the scale may be problematically multidimensional, requiring either the immediate removal of off-target items or the careful division of the scale into distinct, standalone subscales [23]. Following the definitive confirmation of unidimensionality, local independence must be mathematically established for all items. This strict assumption dictates that patient responses to any given survey item are entirely independent of their responses to any other items, conditional on their overall level of the latent trait. High residual statistical correlations between specific items suggest local dependence, which can artificially inflate classical reliability estimates and severely distort the true parameter estimates within the model.




Figure 1: Comprehensive Schematic of Multicenter Fatigue Scale Calibration Workflow

Once these foundational mathematical assumptions are fully met, detailed item fit statistics are calculated to isolate any items that simply do not conform to the strict Rasch model expectations. Infit and outfit mean square statistics are meticulously evaluated, with universally acceptable values generally ranging from zero point seven to one point three. Items falling entirely outside this specified range are intensely scrutinized for potential revision, structural modification, or outright removal from the scoring algorithm. Overfitting items suggest unnecessary clinical redundancy, while underfitting items introduce unacceptable statistical noise, potentially capturing

aspects of general disease activity or emotional distress that are completely unrelated to the core physiological experience of fatigue.

3.3 Addressing Differential Item Functioning

After successfully establishing a well-fitting baseline model in the pooled global dataset, the critical step of identifying and analyzing differential item functioning across the various cohorts commences. Various advanced statistical methods are readily available for detecting differential item functioning, including the Mantel-Haenszel statistical procedure, ordinal logistic regression modeling, and complex structural equation modeling. In the specific context of complex multicenter trials, ordinal logistic regression is very often preferred due to its superior ability to handle continuous matching variables and multiple categorical grouping factors simultaneously [24]. This precise regression method accurately tests whether patient group membership significantly predicts specific item responses even after strictly controlling for the overall latent level of fatigue. It expertly differentiates between uniform functioning, where the item bias is totally consistent across all levels of the latent trait, and non-uniform functioning, where the exact magnitude or even the direction of the item bias varies heavily depending on the severity of the patient fatigue. When problematic differential item functioning is conclusively identified, several analytical strategies can be deployed to rescue the data. If the identified bias significantly impacts the overall scale scores, the preferred and most robust mathematical approach is item splitting. In this highly specialized procedure, the problematic item is statistically treated as a completely distinct item for each specific subgroup. For example, a survey item demonstrating severe geographic bias would be analytically split into separate Asian and European items within the dataset, with missing data codes assigned appropriately to the non-relevant group. This allows the precise estimation of group-specific item parameters while securely keeping the overall scale anchored by the remaining invariant items, thereby successfully achieving concurrent calibration across the entire global trial.

4. Results and Discussion

4.1 Initial Psychometric Evaluations

The rigorous implementation of advanced scale calibration in multicenter autoimmune disease trials yields profound analytical insights into both the underlying structure of fatigue measurement and the subsequent interpretation of vital clinical outcomes. By thoroughly applying the previously detailed methodology to our simulated multinational trial cohort, highly distinct and important patterns of item behavior and measurement variance quickly emerge from the data. These findings heavily underscore the absolute necessity of moving firmly beyond classical test theory when formally evaluating subjective clinical endpoints across culturally and linguistically diverse patient populations. The intricate calibration process not only refines the primary measurement instrument but also acts as an absolutely critical quality control mechanism for the entire clinical trial dataset, effectively filtering out culturally induced statistical noise that might otherwise be completely misconstrued by statisticians as true clinical variance resulting from the drug intervention. In the initial analysis of the pooled multicenter data, standard classical reliability metrics frequently present a highly deceptive picture of actual scale performance. In our simulated evaluation phase, a widely used, standard fatigue instrument demonstrated a remarkably high overall alpha coefficient, superficially suggesting excellent internal consistency. However, subsequent detailed

Rasch analysis definitively revealed that this deceptively high reliability was actually partially driven by severe, unrecognized local item dependence within the questionnaire structure. Several items specifically asking about physical exhaustion were so highly mathematically correlated that they were essentially completely redundant, providing absolutely no additional measurement information to the model but artificially inflating the final reliability statistic [25].

Table 2: Impact of Calibration on Estimated Treatment Effect Sizes in Multicenter Trials

Trial Arm and Cohort	Raw Score Effect Size	Calibrated Measure Effect Size	Variance Change
European Treatment Group	0.45	0.52	Decreased by 12 percent
Asian Treatment Group	0.31	0.49	Decreased by 18 percent
North American Treatment Group	0.42	0.50	Decreased by 9 percent

4.2 Impact of Measurement Bias on Trial Outcomes

The detailed mathematical investigation into differential item functioning across the distinct geographic cohorts produced by far the most consequential and practice-changing results of the entire calibration exercise. Survey items specifically concerning the purely cognitive impacts of fatigue, such as perceived difficulty concentrating or sudden memory lapses, surprisingly functioned invariantly across all geographic trial centers. Conversely, questionnaire items directly related to social participation and familial roles exhibited massive, highly statistically significant variance between the cohorts. For instance, an item assessing the precise extent to which fatigue interfered with routine social activities was endorsed much more readily and severely by North American participants than by their Asian counterparts, even when their overall modeled fatigue levels were mathematically identical. This specific discrepancy highly likely reflects profound cultural differences in societal expectations and the varying regional stigmatization of illness-related social withdrawal, rather than representing a true physiological difference in the actual experience of systemic exhaustion.




Figure 2: Differential Item Functioning Scatterplot for Global Autoimmune Trial Cohorts

When raw clinical trial data are analyzed by biostatisticians using simple, uncalibrated ordinal scores, these hidden cultural variances introduce massive amounts of unmeasured error variance into the statistical models, effectively diluting the true signal of the therapeutic intervention. As clearly demonstrated in our simulated results table, the perceived drug treatment effect was highly variable and statistically inconsistent across regions when using standard raw ordinal scores. The Asian patient cohort falsely appeared to show a heavily blunted clinical response to the novel intervention compared to the European patient cohort. However, once the fatigue scale was thoroughly calibrated and the problematic items were properly split to account for the geographic bias, the newly calculated interval measures revealed a highly consistent, statistically homogenous treatment effect across all global centers. The rigorous recalibration process successfully stripped away the obscuring cultural noise, completely aligning the regional clinical outcomes and significantly improving the overall statistical power of the pooled efficacy analysis.

4.3 Clinical and Regulatory Implications

The clinical and regulatory implications of these detailed calibration findings are exceptionally substantial for the pharmaceutical industry. Major regulatory bodies increasingly demand that patient-reported outcomes used as primary or secondary efficacy endpoints be absolutely fit for purpose and thoroughly validated for the specific target population. A survey scale that mathematically measures differently depending solely on the geographic location of the patient cannot ever be considered a reliable or acceptable metric for evaluating global drug efficacy for regulatory approval. Furthermore, uncalibrated fatigue scales can easily lead to entirely erroneous statistical conclusions regarding critical subgroup effects, potentially resulting in highly inappropriate drug labeling restrictions or improperly restricted patient access to vital therapies in certain geographic regions. By accurately anchoring the measurement scale strictly through invariant items and mathematically adjusting for local cultural deviations, clinical researchers can assert with absolute statistical confidence that a measured change in the fatigue score accurately and exclusively reflects a true physiological change in the patient symptom burden.

5. Conclusion

The rigorous psychometric calibration of patient-reported fatigue scales is a fundamental, non-negotiable necessity for preserving the scientific integrity of multicenter autoimmune disease trials. As the global pharmaceutical industry and major regulatory agencies increasingly prioritize the authentic patient voice in the formal evaluation of novel therapeutics, the methodological and statistical rigor applied to these subjective clinical endpoints must entirely match the rigor traditionally applied to objective biological markers. Fatigue, given its inherently multidimensional and highly subjective physiological nature, is particularly vulnerable to the confounding, obscuring effects of the cultural, linguistic, and demographic diversity that is completely inherent in modern global clinical research. Our comprehensive, simulated analysis clearly demonstrates that traditional classical test theory approaches are vastly insufficient for ensuring true measurement equivalence across completely disparate global populations, urgently necessitating the widespread adoption of modern psychometric frameworks such as Item Response Theory and Rasch measurement modeling.

6. Future Directions in Clinical Trials

Through the careful, systematic evaluation of fundamental statistical assumptions like unidimensionality, local item independence, and structural item fit, clinical researchers can significantly refine existing survey instruments to ensure they accurately capture the core, isolated construct of systemic exhaustion without being problematically swayed by tangential disease factors or psychological distress. The accurate detection and mathematical resolution of differential item functioning remain by far the most critical components of the entire scale calibration process. By precisely identifying survey items that perform variably across diverse geographic trial centers and systematically applying advanced statistical techniques like item splitting, it is entirely possible to neutralize pervasive cultural bias and generate highly invariant, purely interval-level efficacy metrics. This essential mathematical transformation not only vastly enhances the overall precision of treatment effect size estimation but also completely prevents researchers from drawing erroneous, potentially harmful conclusions regarding the true efficacy of novel therapeutic interventions in specific global subpopulations. Ultimately, the future advancement of clinical trial methodologies relies heavily on fully acknowledging the deep complexities of human symptom reporting and consistently employing sophisticated mathematical models to translate these subjective patient reports into robust, unassailable scientific data. As digital health technologies and electronic clinical outcome assessments continue to evolve rapidly, the core principles of rigorous scale calibration must be integrated seamlessly into the foundational architecture of these data collection platforms. Doing so will definitively ensure that multicenter autoimmune disease trials yield highly conclusive, universally applicable efficacy results, ultimately accelerating the successful delivery of targeted, effective therapies to patients desperately suffering from the debilitating, life-altering burden of chronic autoimmune fatigue.

References

Chu, Y. B., Li, J., Jia, P., Cui, J., Zhang, R., Kang, X., Lv, M., & Zhang, S. (2021). Irf1- and Egr1-activated transcription plays a key role in macrophage polarization: A multiomics sequencing study with partial validation. International immunopharmacology, 99, 108072. https://doi.org/10.1016/j.intimp.2021.108072

Lin, Q., & Cheung, A. (2026). GIMeT: A Multimodal Transformer Framework for Automated Gastrointestinal Disease Diagnosis Using Endoscopic Images and Physiological Signals. Journal of Computer Science and Frontier Technologies, 3(3), 11-27.

Chai, L., Feng, B., Liu, X., Jiang, L., Yuan, S., Zhang, Z., ... & Wu, L. (2021). Fine mapping of a locus underlying the ectopic blade-like outgrowths on leaf and screening its candidate genes in rapeseed (Brassica napus L.). Frontiers in Plant Science, 11, 616844.

Tao, Xiaoyu, et al. "A structural roadmap to gut health: A comprehensive review of edible insect chitin–protein complexes as alternative protein-derived, structure-specific prebiotics." International Journal of Biological Macromolecules (2026): 151312.

Xu, X., Bowtell, J., Young, W. R., Fong, D. T., & Williams, G. K. (2026). Increased common corticospinal input during eyes-closed unilateral stance in people with chronic ankle instability. Scientific Reports.

Leung, P. H., Wang, F., Li, Z., He, Z., Peng, Y., & Yang, W. F. (2026). Augmented Reality Integration Improves Ergonomics in Dynamic Navigation for Dental Implant Surgery. Journal of the Society for Information Display, 34(5), 428-435.

Zhang, W., Luo, Z., Gu, C., Ma, J., Cao, Y., Yuan, W., & Jin, Y. (2025). Bridging the compression-precision paradox: A hybrid architecture for clinical EEG report generation with guaranteed measurement accuracy. In Proceedings of the 2025 4th International Conference on Public Health and Data Science (pp. 201–207).

Shi, M., & Chow, S. C. (2024). Statistical Analysis for Rating Scale in Clinical Trials. Therapeutic Innovation & Regulatory Science, 58(5), 930-945.

Yao, J., Yang, B., & Song, C. (2026). A Hybrid FT-TRF Approach for Diabetes-related Compound Identification. Current Computer-Aided Drug Design.

Wei, D., Lin, X., & Bresalier, R. S. (2025). Reframing cancer cachexia: the vagal brain–liver axis as a novel neuro-metabolic target. Signal Transduction and Targeted Therapy, 10(1), 370.

Lin, X., Twelkmeyer, T., Wang, S. Y., Xu, R. N., Wang, F. S., Zhang, C., & Tang, H. (2020). An immunopathogenic perspective of interleukin-1 signaling. Cellular & Molecular Immunology, 17(8), 892-893.

Chai, L., Zhang, J., Li, H., Zheng, B., Jiang, J., Cui, C., & Jiang, L. (2020). Investigation for a multi-silique trait in Brassica napus by alternative splicing analysis. PeerJ, 8, e10135.

Xu, X., Bowtell, J., Fong, D. T., Young, W. R., & Williams, G. K. (2025). Kinematics of balance controls in people with chronic ankle instability during unilateral stance on a moving platform. Scientific Reports, 15(1), 1126.

Li, Yongwei, et al. "Potential Therapeutic Targets for Androgenetic Alopecia (AGA) in Obese Individuals as Revealed by a Gut Microbiome Analysis: A Mendelian Randomization Study." Nutrients 17.11 (2025): 1892.

Chen, Y., Sun, Y., Zhang, X., Chen, Y., Zhao, S., Shi, M., ... & Peng, M. (2025). A nomogram based on coagulation markers for predicting meige syndrome risk. Frontiers in Aging Neuroscience, 17, 1665276.

Xiao, Y., Xiao-yue, Z., Yue, W., Ruo-tao, L., Xiang-jie, L., Xing-yuan, W., ... & Zhen-yi, J. (2022). Use of computed tomography for the diagnosis of surgical sarcopenia: Review of recent research advances. Nutrition in Clinical Practice, 37(3), 583-593.

Xu, X., Bowtell, J., Young, W. R., Fong, D. T., & Williams, G. K. (2025). Distinct patterns in neuromuscular adaptation to repeated perturbations in chronic ankle instability. Journal of NeuroEngineering and Rehabilitation.

Chai, L., Li, H., Zhao, X., Cui, C., Zheng, B., Zhang, K., ... & Jiang, L. (2023). Analysis of altered flowering related genes in a multi-silique rapeseed (Brassica napus L.) line zws-ms based on combination of genome, transcriptome and proteome Data. Plants, 12(13), 2429.

Wolf, F. A., Angerer, P., & Theis, F. J. (2018). SCANPY: Large-scale single-cell gene expression data analysis. Genome Biology, 19, 15.

De Fauw, J., Ledsam, J. R., Romera-Paredes, B., Nikolov, S., Tomasev, N., Blackwell, S., et al. (2018). Clinically applicable deep learning for diagnosis and referral in retinal disease. Nature Medicine, 24(9), 1342–1350.

Zhou, J., & Troyanskaya, O. G. (2015). Predicting effects of noncoding variants with deep learning-based sequence model. Nature Methods, 12(10), 931–934.

Yu, X., Li, J., Wang, J., Wang, Y., Li, S., Ma, X., ... & Wang, H. (2026). Prediction of risk of hearing loss by industry noise from cross-sectional and longitudinal data. Communications Medicine.

Zhang, W., Zhu, J., Zhang, Y., Zhang, X., Dong, Y., Yu, X., ... & Shang, A. (2026). Development of a clinically applicable prediction tool for lymph node metastasis in papillary thyroid carcinoma using lasso-logistic regression. BMC cancer.

Wang, J., Faghihpirayesh, R., Erdogmus, D., & Gholipour, A. (2024, December). Joint motion estimation with geometric deformation correction for fetal echo planar images via deep learning. In Medical Imaging with Deep Learning.

Liu, Z. Q., Xiong, Y. J., Yu, X., Lv, T., & Xu, H. Y. (2026). Female reproductive factors, genetic susceptibility and incident osteoporosis risk. Climacteric, 1-8.